

Clinical Development - Counter-Terrorism Vaccines

Center for Biologics
Evaluation and Research

October 23, 2003
Jeff Brady, M.D., M.P.H.
Medical Officer, CBER, DVRPA



U.S. Department of Health and Human Services

Food and Drug Administration

Purpose of Presentation

- Overview of preventive vaccine clinical development
- Focus on Phase 1 and 2 trials
- Identify special considerations for vaccine development
- Encourage sponsors to identify global development goals early
 - target populations
 - label indications
 - anticipated use

Bio-Terrorism Diseases / Agents

Category A

- Smallpox
- Anthrax
- Botulinum toxin
- Plague
- Tularemia
- Viral hemorrhagic fevers (Ebola, Marburg)
- Arenaviruses (Lassa, Junin)

Category C

- Nipah virus
- Hantavirus
- Yellow Fever
- Multidrug-resistant tuberculosis
- Tickborne hemorrhagic fever viruses
- Tickborne encephalitis viruses

Category B

- Brucellosis
- Glanders
- Q-fever
- Alphaviruses (EEE, WEE, VEE)
- Epsilon toxin (*Cl. perfringens*)
- Ricin toxin
- Staphylococcal enterotoxin B
- Salmonella spp.
- Cholera
- *E. coli* 0157:H7
- Cryptosporidiosis
- Shigellosis

Vaccine Development

Pre-IND

IND

Development
of Rationale
Based on
Disease
Pathogenesis



Immunogen
Identification




Development
of
Manufacturing
Process;
Preclinical &
Nonclinical
Studies;
Protocol
Concept Sheet

Clinical
Studies;
Additional
Non-clinical
Preclinical
Work;
Scale-up

IND = Investigational New Drug application


Stages of Review and Regulation



IND = Investigational New Drug Application;
BLA= Biologics License Application

Recommended Meetings with FDA

(21 CFR 312.47)




IND =Investigational New Drug Application
BLA =Biologics License Application

*Shouldn't be a surprise

Stages of Review and Regulation

Clinical Investigational Plan



IND = Investigational New Drug Application;
BLA= Biologics License Application

Phase 1 Study

General Considerations

- **Objectives and endpoints**
 - **Primary: Safety and tolerability**
 - **Secondary: Preliminary immunogenicity**
- **Closely monitored (safety)**
- **Adults, at least for first phase 1 study**
- **Sample Size**
 - **Small study: e.g., 20 to 80**
- **Special instructions for vaccinees, if needed**

Phase 1 Study

Features and Components

- Consider vaccine-specific features when planning trial (e.g., live vaccine)
- Develop Inclusion and Exclusion Criteria
 - Healthy adult volunteers
 - Age range: 18-40 years recommended (esp. for first phase 1 study)
 - Special considerations
 - age, serostatus, concomitant medications allowed, etc.
 - where applicable, vaccinee contacts
 - E.g., vaccinia

Safety Monitoring

- **Goals:**
 - Protect subjects by monitoring local, systemic, and potential end-organ toxicity
 - Identify major toxicity
- **Clinic visits**
 - Symptom review, diary cards
 - Clinical exam
- **Laboratory studies**
 - CBC: hematologic
 - Chemistries: e.g., hepatic, renal (U/A), endocrine
 - Others? Per pre-clinical toxicology study, previous experience with similar vaccines, etc.

Safety Monitoring (cont'd.)

- **Safety and activity (e.g., immunogenicity):**
 - **Items to be assessed/time schedule**
(Well organized summary in a table)
 - **Active post-vaccination monitoring**
 - **Monitoring tools**
 - **Submit to IND with protocol, regardless of Phase**
 - **Prototype Case Report Forms (CRFs)**
 - **Diary cards**
 - **Scripted interviews**
 - **Other, e.g., photographs**

Safety Monitoring (cont'd.)


- **Toxicity Grading Scales**
 - Define grades for specifically monitored parameters (clinical and laboratory AEs)
 - Based on healthy volunteers
- **Stopping rules**
 - Provide specific criteria
 - Address grade 3 (severe) or grade 4 (serious) adverse events
 - If criteria met, stop vaccination and investigate
 - Safety review
 - If appropriate, resume study +/- changes to protocol / I.C.

Phase 1 Study

Features and Components (cont'd.)

- **Dose escalation**
 - **Even in first Phase 1 study**
 - **Provide details of dose escalation scheme**
 - **Clear criteria for dose escalation**
 - **Safety review of lowest dose cohort**

Stages of Review and Regulation



IND = Investigational New Drug Application;
BLA= Biologics License Application

Phase 2 Study

General Considerations

- **Goals:**
 - **Immunogenicity**
 - **Dose-ranging data**
 - **Identify preferred dose, schedule, formulation, route of administration for advancement to Phase 3**
 - **Safety**
 - **More precise estimates of common adverse events**
 - **Local reactogenicity**
 - **Systemic effects**


Phase 2 Clinical Trials

- Up to several hundred subjects in a trial
- Broader study population
- Often randomized & controlled
- Vaccine-elicited immune responses
 - Qualitative
 - Quantitative
 - Duration
- Safety
- Pilot evaluation of efficacy endpoints (where feasible)

Phase 2 Clinical Trials

- **Planning for Phase 3**
- **Logistics and Protocol:**
 - **Compliance with protocol**
 - **Accrual of subjects**
 - **Target populations for licensure**
 - **Monitoring tools**
 - **Sample handling**

Stages of Review and Regulation



IND = Investigational New Drug Application;
BLA= Biologics License Application

Phase 3 Development General Considerations

- **Develop adequate safety, immunogenicity, and efficacy data to support**
 - **Proposed use(s) and indication(s)**
 - **Target population(s)**

Phase 3 Study

General Considerations

- Objectives and Endpoints:
 - Pivotal efficacy - options
 - 1) Clinical endpoint, if feasible
 - 2) Immune response endpoint
 - 3) “Animal Rule”, if appropriate
 - Pivotal pre-licensure safety database
 - Sample Size: Thousands for safety in humans, regardless of path to licensure

Phase 3 Vaccine Efficacy Trial Protocol

- Study population/background epidemiology
- Control group
- Randomization scheme/Study masking
- Items assessed/time schedule:
 - Clinical & lab parameters: safety, immunogenicity, microbiology and efficacy
- Prospective 1^o & 2^o efficacy endpoints

Efficacy Trial Endpoints

- Clinical relevance of case definition, esp. for primary endpoint
- Specificity of case definition emphasized*
- Validation of assays before efficacy study
 - Performance Parameters

*Lachenbruch PA: Sensitivity, Specificity, & Vaccine Efficacy. Controlled Clin Trials 19:569-574, 1998.


*Orenstein WA et al. Assessing Vaccine Efficacy in the Field: Further Observations. Epidemiol Rev 10: 212-241, 1988.

Phase 3 Protocols

- **Use of animal rule**
 - **Criteria for dose used in Phase 3 must consider results of animal efficacy studies**
 - **Compare immune responses in animals and humans**
- **Develop Phase 3 safety data at appropriate dose**
 - **Randomized, controlled safety data most interpretable**
 - **Appropriate control group**

Stages of Review and Regulation

Clinical Investigational Plan



IND = Investigational New Drug Application;
BLA= Biologics License Application

Post-marketing Studies

- **Limitations of pre-licensure studies**
 - Rare adverse events
 - Delayed onset / long term effects
 - Sub-population
 - Efficacy
- **Specific post-marketing commitments at the time of approval**
 - Review of recent vaccine approval letters may be instructive

Published Guidance

FDA, ICH

FDA Guidance Documents for Industry

- <http://www.fda.gov/cber/guidelines.htm>
- <http://www.fda.gov/cder/guidance/>

International Conference on Harmonisation

- E6: <http://www.ich.org/ich5e.html#GCP>

Conclusions: Counter-terrorism vaccine development

- **Early and frequent regulatory communication**
 - **Pre-IND Meeting: feedback on phase 1 trial design**
 - **Early articulation of development goals**
 - **Target population(s)**
 - **Indication(s)**

Conclusions: Counter-terrorism vaccine development (cont'd.)

- “Animal rule” if applicable
- Develop data on relevant dose in Phase 2 to be investigated in Phase 3
- Adequate safety data

Acknowledgements

- **Division of Vaccines and Related Products Applications (DVRPA)**
- **Dr. Karen L. Goldenthal**
- **Vaccine Clinical Trials Branch (VCTB)**
- **Dr. Antonia Geber**
- **Dr. Douglas Pratt**
- **Dr. Joe Toerner**
- **Dr. Steve Rosenthal**